

# On polynomials for the calculation of sums of powers of successive integers and Bernoulli numbers deduced from the Pascal's triangle

by *Giorgio Pietrocola*

[giorgio.pietrocola\[at\]gmail.com](mailto:giorgio.pietrocola[at]gmail.com)

June 2017

(updating 14.9.2017 : 4.1, 4.2)

(updating 1.10.2017: 5.0,6.0)

(updating 23.6.2020 7.0)

Abstract: In the "twin" theorems [1A](#) and [1B](#), two matrices derived from the Pascal's triangle, the first of which has alternate signs, are taken into account, and it is shown that their inverse gives, with a minimum variation in the second case, the coefficients of famous polynomials for calculating the sums of powers of successive integers (fig.3). These coefficients are determined here without the help of Bernoulli's numbers. In the theorems [2A](#) and [2B](#), these famous numbers are obtained starting from other matrices closely related to the previous ones.

**1.Theorem 1A. It will be shown that the polynomial coefficients for calculating the sum of powers of n successive integers can be obtained by inverting a particular alternate-sign matrix in relationship to Pascal's triangle.**

Suppose  $A_m$  is a square matrix of order m defined as:

$$a_{j,k} = \begin{cases} 0 & \text{if } k > j \\ \binom{j}{k-1} * (-1)^{j+k} & \text{otherwise} \end{cases}$$

This is a triangular matrix where the Pascal's triangle appears, with alternate signs, private of the last element of each line. Its main diagonal is therefore formed by the succession of the positive integers whose product (m!) gives the determinant.

Therefore:

$$\begin{aligned}
 A_m * \begin{pmatrix} \sum_{i=1}^n i^0 \\ \sum_{i=1}^n i^1 \\ \sum_{i=1}^n i^2 \\ \dots \\ \sum_{i=1}^n i^{m-1} \end{pmatrix} &= A_m * \sum_{i=1}^n \begin{pmatrix} i^0 \\ i^1 \\ i^2 \\ \dots \\ i^{m-1} \end{pmatrix} = \sum_{i=1}^n A_m * \begin{pmatrix} i^0 \\ i^1 \\ i^2 \\ \dots \\ i^{m-1} \end{pmatrix} = \\
 &= \sum_{i=1}^n \begin{pmatrix} i^1 - (i-1)^1 \\ i^2 - (i-1)^2 \\ i^3 - (i-1)^3 \\ \dots \\ i^m - (i-1)^m \end{pmatrix} = \begin{pmatrix} n^1 \\ n^2 \\ n^3 \\ \dots \\ n^m \end{pmatrix}
 \end{aligned}$$

In the first two steps we applied the distributive property, in the third step multiplying the  $A_m$  matrix by the vectors of powers, we get:

$$\begin{aligned}
 \sum_{k=1}^m a_{j,k} * i^{k-1} &= \sum_{k=1}^j \binom{j}{k-1} * (-1)^{j+k} * i^{k-1} = \\
 &= \sum_{k=1}^j (-1)^{2k-1} \binom{j}{k-1} * (-1)^{j-k+1} * i^{k-1} = \\
 &= - \sum_{k=1}^j \binom{j}{k-1} * (-1)^{j-k+1} * i^{k-1} = \\
 &= -(-i^j + \sum_{k=1}^{j+1} \binom{j}{k-1} * (-1)^{j-k+1} * i^{k-1}) = \\
 &= -(-i^j + (i-1)^j) = i^j - (i-1)^j
 \end{aligned}$$

Therefore we find:

$$A_m * \begin{pmatrix} \sum_{i=1}^n i^0 \\ \sum_{i=1}^n i^1 \\ \sum_{i=1}^n i^2 \\ \dots \\ \sum_{i=1}^n i^{m-1} \end{pmatrix} = \begin{pmatrix} n \\ n^2 \\ n^3 \\ \dots \\ n^m \end{pmatrix}$$

Multiplying the two left hand members by the inverse matrix of  $A_m$  we get:

$$\begin{pmatrix} \sum_{i=1}^n i^0 \\ \sum_{i=1}^n i^1 \\ \sum_{i=1}^n i^2 \\ \dots \\ \sum_{i=1}^n i^{m-1} \end{pmatrix} = A_m^{-1} \begin{pmatrix} n \\ n^2 \\ n^3 \\ \dots \\ n^m \end{pmatrix}$$

Thus, the theorem provides the famous polynomials for computing the sum of powers of successive integers already studied by Faulhaber in the seventeenth century

Example given m=11

$$A_{11} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -1 & 2 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & -3 & 3 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -1 & 4 & -6 & 4 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & -5 & 10 & -10 & 5 & 0 & 0 & 0 & 0 & 0 & 0 \\ -1 & 6 & -15 & 20 & -15 & 6 & 0 & 0 & 0 & 0 & 0 \\ 1 & -7 & 21 & -35 & 35 & -21 & 7 & 0 & 0 & 0 & 0 \\ -1 & 8 & -28 & 56 & -70 & 56 & -28 & 8 & 0 & 0 & 0 \\ 1 & -9 & 36 & -84 & 126 & -126 & 84 & -36 & 9 & 0 & 0 \\ -1 & 10 & -45 & 120 & -210 & 252 & -210 & 120 & -45 & 10 & 0 \\ 1 & -11 & 55 & -165 & 330 & -462 & 462 & -330 & 165 & -55 & 11 \end{pmatrix}$$

And applying the theorem:

$$\begin{pmatrix} \sum_{i=1}^n i^0 \\ \sum_{i=1}^n i^1 \\ \sum_{i=1}^n i^2 \\ \sum_{i=1}^n i^3 \\ \sum_{i=1}^n i^4 \\ \sum_{i=1}^n i^5 \\ \sum_{i=1}^n i^6 \\ \sum_{i=1}^n i^7 \\ \sum_{i=1}^n i^8 \\ \sum_{i=1}^n i^9 \\ \sum_{i=1}^n i^{10} \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ \frac{1}{2} & \frac{1}{2} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ \frac{1}{6} & \frac{1}{2} & \frac{1}{3} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & \frac{1}{4} & \frac{1}{2} & \frac{1}{4} & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -\frac{1}{30} & 0 & \frac{1}{3} & \frac{1}{2} & \frac{1}{5} & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -\frac{1}{12} & 0 & \frac{5}{12} & \frac{1}{2} & \frac{1}{6} & 0 & 0 & 0 & 0 & 0 \\ \frac{1}{42} & 0 & -\frac{1}{6} & 0 & \frac{1}{2} & \frac{1}{2} & \frac{1}{7} & 0 & 0 & 0 & 0 \\ 0 & \frac{1}{12} & 0 & -\frac{7}{24} & 0 & \frac{7}{12} & \frac{1}{2} & \frac{1}{8} & 0 & 0 & 0 \\ -\frac{1}{30} & 0 & \frac{2}{9} & 0 & -\frac{7}{15} & 0 & \frac{2}{3} & \frac{1}{2} & \frac{1}{9} & 0 & 0 \\ 0 & -\frac{3}{20} & 0 & \frac{1}{2} & 0 & -\frac{7}{10} & 0 & \frac{3}{4} & \frac{1}{2} & \frac{1}{10} & 0 \\ \frac{5}{66} & 0 & -\frac{1}{2} & 0 & 1 & 0 & -1 & 0 & \frac{5}{6} & \frac{1}{2} & \frac{1}{11} \end{pmatrix} * \begin{pmatrix} n \\ n^2 \\ n^3 \\ n^4 \\ n^5 \\ n^6 \\ n^7 \\ n^8 \\ n^9 \\ n^{10} \\ n^{11} \end{pmatrix}$$

To note in the first column of the square matrix the appearance of Bernoulli's original numbers.

Or likewise:

$$\begin{aligned}
\sum_{i=1}^n i^0 &= n \\
\sum_{i=1}^n i^1 &= \frac{1}{2}n + \frac{1}{2}n^2 \\
\sum_{i=1}^n i^2 &= \frac{1}{6}n + \frac{1}{2}n^2 + \frac{1}{3}n^3 \\
\sum_{i=1}^n i^3 &= \frac{1}{4}n^2 + \frac{1}{2}n^3 + \frac{1}{4}n^4 \\
\sum_{i=1}^n i^4 &= -\frac{1}{30}n + \frac{1}{3}n^3 + \frac{1}{2}n^4 + \frac{1}{5}n^5 \\
\sum_{i=1}^n i^5 &= -\frac{1}{12}n^2 + \frac{5}{12}n^4 + \frac{1}{2}n^5 + \frac{1}{6}n^6 \\
\sum_{i=1}^n i^6 &= \frac{1}{42}n - \frac{1}{6}n^3 + \frac{1}{2}n^5 + \frac{1}{2}n^6 + \frac{1}{7}n^7 \\
\sum_{i=1}^n i^7 &= \frac{1}{12}n^2 - \frac{7}{24}n^4 + \frac{7}{12}n^6 + \frac{1}{2}n^7 + \frac{1}{8}n^8 \\
\sum_{i=1}^n i^8 &= -\frac{1}{30}n + \frac{2}{9}n^3 - \frac{7}{15}n^4 + \frac{2}{3}n^6 + \frac{1}{2}n^8 + \frac{1}{9}n^9 \\
\sum_{i=1}^n i^9 &= -\frac{3}{20}n^2 + \frac{1}{2}n^4 - \frac{7}{10}n^6 + \frac{3}{4}n^8 + \frac{1}{2}n^9 + \frac{1}{10}n^{10} \\
\sum_{i=1}^n i^{10} &= \frac{5}{66}n - \frac{1}{2}n^3 + n^5 - n^7 + \frac{5}{6}n^9 + \frac{1}{2}n^{10} + \frac{1}{11}n^{11}
\end{aligned}$$

These are the polynomials studied by Faulhaber and Bernoulli

... Atque si porrò ad altiores gradatim potestates pergere, levique negotio sequentem adornare laterculum licet :

*Summae Potestatum*

$$f n = \frac{1}{2}nn + \frac{1}{2}n$$

$$f nn = \frac{1}{3}n^3 + \frac{1}{2}nn + \frac{1}{6}n$$

$$f n^3 = \frac{1}{4}n^4 + \frac{1}{2}n^3 + \frac{1}{4}nn$$

$$f n^4 = \frac{1}{5}n^5 + \frac{1}{2}n^4 + \frac{1}{3}n^3 - \frac{1}{30}n$$

$$f n^5 = \frac{1}{6}n^6 + \frac{1}{2}n^5 + \frac{5}{12}n^4 - \frac{1}{12}nn$$

$$f n^6 = \frac{1}{7}n^7 + \frac{1}{2}n^6 + \frac{1}{2}n^5 - \frac{1}{6}n^3 + \frac{1}{42}n$$

$$f n^7 = \frac{1}{8}n^8 + \frac{1}{2}n^7 + \frac{7}{12}n^6 - \frac{7}{24}n^4 + \frac{1}{12}nn$$

$$f n^8 = \frac{1}{9}n^9 + \frac{1}{2}n^8 + \frac{2}{3}n^7 - \frac{7}{15}n^5 + \frac{2}{9}n^3 - \frac{1}{30}n$$

$$f n^9 = \frac{1}{10}n^{10} + \frac{1}{2}n^9 + \frac{3}{4}n^8 - \frac{7}{10}n^6 + \frac{1}{2}n^4 - \frac{1}{12}nn$$

$$f n^{10} = \frac{1}{11}n^{11} + \frac{1}{2}n^{10} + \frac{5}{6}n^9 - 1n^7 + 1n^5 - \frac{1}{2}n^3 + \frac{5}{66}n$$

Quin imò qui legem progressionis inibi attentuis ensperexit, eundem etiam continuare poterit absque his ratiociniorum ambabimus : Sumtâ enim c pro potestatis cujuslibet exponente, fit summa omnium  $n^c$  seu

$$\begin{aligned} \int n^c &= \frac{1}{c+1}n^{c+1} + \frac{1}{2}n^c + \frac{c}{2}An^{c-1} + \frac{c \cdot c - 1 \cdot c - 2}{2 \cdot 3 \cdot 4}Bn^{c-3} \\ &+ \frac{c \cdot c - 1 \cdot c - 2 \cdot c - 3 \cdot c - 4}{2 \cdot 3 \cdot 4 \cdot 5 \cdot 6}Cn^{c-5} \\ &+ \frac{c \cdot c - 1 \cdot c - 2 \cdot c - 3 \cdot c - 4 \cdot c - 5 \cdot c - 6}{2 \cdot 3 \cdot 4 \cdot 5 \cdot 6 \cdot 7 \cdot 8}Dn^{c-7} \dots \& \text{ ita deinceps,} \end{aligned}$$

exponentem potestatis ipsius n continué minuendo binario, quosque perveniatur ad n vel nn. Literae capitales A, B, C, D & c. ordine denotant coëfficientes ultimorum terminorum pro  $f nn$ ,  $f n^4$ ,  $f n^6$ ,  $f n^8$ , & c. nempe

$$A = \frac{1}{6}, B = -\frac{1}{30}, C = \frac{1}{42}, D = -\frac{1}{30}.$$

*fig.3: Ars Conjectandi di Jakob Bernoulli (1654-1705) Published in 1713. (the monomial of degree 2 of polinomial of degree 10 is wrong)*

**2.Theorem 2A. It is proved as a corollary of the previous theorem 1A, a formula for obtaining Bernoulli numbers from the Pascal triangle**

We have seen that the matrix  $A_n$  so defined

$$a_{j,k} = \begin{cases} 0 & \text{if } k > j \\ \binom{j}{k-1} * (-1)^{j+k} & \text{otherwise} \end{cases}$$

has an inverse matrix that has the first-degree coefficients of Faulhaber polynomials in the first column. These correspond to "second Bernoulli numbers" or even "Bernoulli's original numbers" where  $B_1 = + \frac{1}{2}$  stands. The most commonly considered "first Bernoulli numbers" whose sequence differs only for  $B_1 = -\frac{1}{2}$ .

Recalling the Laplace algorithm for the calculation of the inverse matrices we have;

$$c_{n,1} = B_{n-1} = (-1)^{n+1} \frac{|A_{1,n}|}{|A_n|} = (-1)^{n+1} \frac{|A_{1,n}|}{n!}$$

where:

$c_{n,1}$  indicates the corresponding element of  $a_{n,1}$  in the reverse matrix

$|A_n| = n!$  Is the determinant of the triangular matrix.

$A_{1,n}$  is the algebraic complement (obtained by deleting first row and last column) relative to the corresponding  $a_{1,n}$  element of  $a_{n,1}$  in the transposition for the inverse calculation.

It is therefore possible to define Bernoulli numbers as follows:

$$B_n = (-1)^n \frac{|A_n|}{(n+1)!}$$

Where however  $|A_n|$  is the determinant of a  $n$  order matrix defined as follows:

$$a_{i,k} = \begin{cases} 0 & \text{if } k > 1 + i \\ (-1)^{i+k+1} \binom{i+1}{k-1} & \text{otherwise} \end{cases}$$

This formula gives second Bernoulli numbers. If the factor  $(-1)^n$ , is omitted, since the only Bernoulli number from zero with odd index is  $B_1$ , we get the first Bernoulli numbers:

$$B_n = \frac{|A_n|}{(n+1)!}$$

Examples:

$$B_1 = \frac{|-1|}{2!} = -\frac{1}{2}$$

$$B_2 = \frac{\begin{vmatrix} -1 & 2 \\ 1 & -3 \end{vmatrix}}{3!} = \frac{1}{6}$$



$$B_3 = \frac{\begin{vmatrix} -1 & 2 & 0 \\ 1 & -3 & 3 \\ -1 & 4 & -6 \end{vmatrix}}{4!} = \frac{0}{24} = 0$$

$$B_4 = \frac{\begin{vmatrix} -1 & 2 & 0 & 0 \\ 1 & -3 & 3 & 0 \\ -1 & 4 & -6 & 4 \\ 1 & -5 & 10 & -10 \end{vmatrix}}{5!} = \frac{-4}{120} = -\frac{1}{30}$$

$$B_5 = \frac{\begin{vmatrix} -1 & 2 & 0 & 0 & 0 \\ 1 & -3 & 3 & 0 & 0 \\ -1 & 4 & -6 & 4 & 0 \\ 1 & -5 & 10 & -10 & 5 \\ -1 & 6 & -15 & 20 & -15 \end{vmatrix}}{6!} = \frac{0}{720} = 0$$

$$B_6 = \frac{\begin{vmatrix} -1 & 2 & 0 & 0 & 0 & 0 \\ 1 & -3 & 3 & 0 & 0 & 0 \\ -1 & 4 & -6 & 4 & 0 & 0 \\ 1 & -5 & 10 & -10 & 5 & 0 \\ -1 & 6 & -15 & 20 & -15 & 6 \\ 1 & -7 & 21 & -35 & 35 & -21 \end{vmatrix}}{7!} = \frac{120}{5040} = \frac{1}{42}$$

**3.Theorem 1B.** It will be shown that the polynomial coefficients for calculating the sum of powers of n-1 successive integers can be obtained by inverting a particular alternate-sign matrix in relationship to Pascal's triangle.

Suppose  $A_m$  is a square matrix of order m defined as:

$$a_{j,k} = \begin{cases} 0 & \text{if } k > j \\ \binom{j}{k-1} & \text{otherwise} \end{cases}$$

This is a triangular matrix where the triangle of Tartaglia appears private of the last element of each line. Its main diagonal is therefore formed by the succession of the positive integers whose product ( $m!$ ) gives the determinant.

Therefore:

$$\begin{aligned} A_m * \begin{pmatrix} 1 + \sum_{i=1}^{n-1} i^0 \\ \sum_{i=1}^{n-1} i^1 \\ \sum_{i=1}^{n-1} i^2 \\ \dots \\ \sum_{i=1}^{n-1} i^{m-1} \end{pmatrix} &= A_m * \begin{pmatrix} 1 \\ 0 \\ 0 \\ \dots \\ 0 \end{pmatrix} + A_m * \sum_{i=1}^n \begin{pmatrix} i^0 \\ i^1 \\ i^2 \\ \dots \\ i^{m-1} \end{pmatrix} = \begin{pmatrix} 1 \\ 1 \\ 1 \\ \dots \\ 1 \end{pmatrix} + \sum_{i=1}^n A_m * \begin{pmatrix} i^0 \\ i^1 \\ i^2 \\ \dots \\ i^{m-1} \end{pmatrix} = \\ &= \begin{pmatrix} 1 \\ 1 \\ 1 \\ \dots \\ 1 \end{pmatrix} + \sum_{i=1}^n \begin{pmatrix} (i+1)^1 - i^1 \\ (i+1)^2 - i^2 \\ (i+1)^3 - i^3 \\ \dots \\ (i+1)^m - i^m \end{pmatrix} = \begin{pmatrix} 1 \\ 1 \\ 1 \\ \dots \\ 1 \end{pmatrix} + \begin{pmatrix} n^1 - 1 \\ n^2 - 1 \\ n^3 - 1 \\ \dots \\ n^m - 1 \end{pmatrix} = \begin{pmatrix} n \\ n^2 \\ n^3 \\ \dots \\ n^m \end{pmatrix} \end{aligned}$$

In the first two steps we applied the distributive property, in the third step multiplying the  $A_m$  matrix by the vectors of powers, we get:

$$\begin{aligned}
\sum_{k=1}^m a_{j,k} * i^{k-1} &= \sum_{k=1}^j \binom{j}{k-1} * i^{k-1} = \\
&= -i^j + \sum_{k=1}^{j+1} \binom{j}{k-1} * i^{k-1} = \\
&= -i^j + (i+1)^j = (i+1)^j - i^j
\end{aligned}$$

Therefore we find:

$$A_m * \begin{pmatrix} 1 + \sum_{i=1}^{n-1} i^0 \\ \sum_{i=1}^{n-1} i^1 \\ \sum_{i=1}^{n-1} i^2 \\ \dots \\ \sum_{i=1}^{n-1} i^{m-1} \end{pmatrix} = \begin{pmatrix} n \\ n^2 \\ n^3 \\ \dots \\ n^m \end{pmatrix}$$

Multiplying the two left hand members of equation by the inverse matrix of  $A_m$  we get:

$$\begin{pmatrix} 1 + \sum_{i=1}^{n-1} i^0 \\ \sum_{i=1}^{n-1} i^1 \\ \sum_{i=1}^{n-1} i^2 \\ \dots \\ \sum_{i=1}^{n-1} i^{m-1} \end{pmatrix} = A_m^{-1} \begin{pmatrix} n \\ n^2 \\ n^3 \\ \dots \\ n^m \end{pmatrix}$$

Thus, the theorem provides the variant note of the polynomials for calculating the sum of  $n-1$  addendums

Example given m=11

$$A_{11} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 2 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 3 & 3 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 4 & 6 & 4 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 5 & 10 & 10 & 5 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 6 & 15 & 20 & 15 & 6 & 0 & 0 & 0 & 0 & 0 \\ 1 & 7 & 21 & 35 & 35 & 21 & 7 & 0 & 0 & 0 & 0 \\ 1 & 8 & 28 & 56 & 70 & 56 & 28 & 8 & 0 & 0 & 0 \\ 1 & 9 & 36 & 84 & 126 & 126 & 84 & 36 & 9 & 0 & 0 \\ 1 & 10 & 45 & 120 & 210 & 252 & 210 & 120 & 45 & 10 & 0 \\ 1 & 11 & 55 & 165 & 330 & 462 & 462 & 330 & 165 & 55 & 11 \end{pmatrix}$$

applying the theorem

$$\begin{pmatrix} 1 + \sum_{i=1}^{n-1} i^0 \\ \sum_{i=1}^{n-1} i^1 \\ \sum_{i=1}^{n-1} i^2 \\ \sum_{i=1}^{n-1} i^3 \\ \sum_{i=1}^{n-1} i^4 \\ \sum_{i=1}^{n-1} i^5 \\ \sum_{i=1}^{n-1} i^6 \\ \sum_{i=1}^{n-1} i^7 \\ \sum_{i=1}^{n-1} i^8 \\ \sum_{i=1}^{n-1} i^9 \\ \sum_{i=1}^{n-1} i^{10} \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -\frac{1}{2} & \frac{1}{2} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ \frac{1}{6} & -\frac{1}{2} & \frac{1}{3} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & \frac{1}{4} & -\frac{1}{2} & \frac{1}{4} & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -\frac{1}{30} & 0 & \frac{1}{3} & -\frac{1}{2} & \frac{1}{5} & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -\frac{1}{12} & 0 & \frac{5}{12} & -\frac{1}{2} & \frac{1}{6} & 0 & 0 & 0 & 0 & 0 \\ \frac{1}{42} & 0 & -\frac{1}{6} & 0 & \frac{1}{2} & -\frac{1}{2} & \frac{1}{7} & 0 & 0 & 0 & 0 \\ 0 & \frac{1}{12} & 0 & -\frac{7}{24} & 0 & \frac{7}{12} & -\frac{1}{2} & \frac{1}{8} & 0 & 0 & 0 \\ -\frac{1}{30} & 0 & \frac{2}{9} & 0 & -\frac{7}{15} & 0 & \frac{2}{3} & -\frac{1}{2} & \frac{1}{9} & 0 & 0 \\ 0 & -\frac{3}{20} & 0 & \frac{1}{2} & 0 & -\frac{7}{10} & 0 & \frac{3}{4} & -\frac{1}{2} & \frac{1}{10} & 0 \\ \frac{5}{66} & 0 & -\frac{1}{2} & 0 & 1 & 0 & -1 & 0 & \frac{5}{6} & -\frac{1}{2} & \frac{1}{11} \end{pmatrix} * \begin{pmatrix} n \\ n^2 \\ n^3 \\ n^4 \\ n^5 \\ n^6 \\ n^7 \\ n^8 \\ n^9 \\ n^{10} \\ n^{11} \end{pmatrix}$$

To note in the first column of the square matrix the appearance of first Bernoulli numbers.

Or likewise:

$$\sum_{i=1}^{n-1} i^0 = n - 1$$

$$\sum_{i=1}^{n-1} i^1 = -\frac{1}{2}n + \frac{1}{2}n^2$$

$$\sum_{i=1}^{n-1} i^2 = \frac{1}{6}n - \frac{1}{2}n^2 + \frac{1}{3}n^3$$

$$\sum_{i=1}^{n-1} i^3 = \frac{1}{4}n^2 - \frac{1}{2}n^3 + \frac{1}{4}n^4$$

$$\sum_{i=1}^{n-1} i^4 = -\frac{1}{30}n + \frac{1}{3}n^3 - \frac{1}{2}n^4 + \frac{1}{5}n^5$$

$$\sum_{i=1}^{n-1} i^5 = -\frac{1}{12}n^2 + \frac{5}{12}n^4 - \frac{1}{2}n^5 + \frac{1}{6}n^6$$

$$\sum_{i=1}^{n-1} i^6 = \frac{1}{42}n - \frac{1}{6}n^3 + \frac{1}{2}n^5 - \frac{1}{2}n^6 + \frac{1}{7}n^7$$

$$\sum_{i=1}^{n-1} i^7 = \frac{1}{12}n^2 - \frac{7}{24}n^4 + \frac{7}{12}n^6 - \frac{1}{2}n^7 + \frac{1}{8}n^8$$

$$\sum_{i=1}^{n-1} i^8 = -\frac{1}{30}n + \frac{2}{9}n^3 - \frac{7}{15}n^4 + \frac{2}{3}n^6 - \frac{1}{2}n^8 + \frac{1}{9}n^9$$

$$\sum_{i=1}^n i^9 = -\frac{3}{20}n^2 + \frac{1}{2}n^4 - \frac{7}{10}n^6 + \frac{3}{4}n^8 - \frac{1}{2}n^9 + \frac{1}{10}n^{10}$$

$$\sum_{i=1}^n i^{10} = \frac{5}{66}n - \frac{1}{2}n^3 + n^5 - n^7 + \frac{5}{6}n^9 - \frac{1}{2}n^{10} + \frac{1}{11}n^{11}$$

Note that these polynomials differ from those originally considered by Faulhaber and Bernoulli only for the sign of the coefficient of the monomy having a degree less than that of the polynomial belonging.

**4. Theorem 2B.** It is proved as a corollary of the previous theorem 1A, a formula for obtaining Bernoulli numbers from the Pascal's triangle

We have seen that the matrix  $A_n$  so defined

$$a_{j,k} = \begin{cases} 0 & \text{if } k > j \\ \binom{j}{k-1} & \text{otherwise} \end{cases}$$

has an inverse matrix that has the first-degree coefficients of polynomials in the first column. These correspond to "first Bernoulli numbers" where  $B_1 = -\frac{1}{2}$  stands. The "second Bernoulli numbers" differs only for  $B_1 = +\frac{1}{2}$ .

Recalling the Laplace algorithm for the calculation of the inverse matrices we have;

$$c_{n,1} = B_{n-1} = (-1)^{n+1} \frac{|A_{1,n}|}{|A_n|} = (-1)^{n+1} \frac{|A_{1,n}|}{n!}$$

where:

$c_{n,1}$  indicates the corresponding element of  $a_{n,1}$  in the reverse matrix

$|A_n| = n!$  Is the determinant of the triangular matrix.

$A_{1,n}$  is the algebraic complement (obtained by deleting first row and last column) relative to the corresponding  $a_{1,n}$  element of  $a_{n,1}$  in the transposition for the inverse calculation.

It is therefore possible to define Bernoulli numbers as follows:

$$B_n = (-1)^n \frac{|A_n|}{(n+1)!}$$

Where however  $|A_n|$  is the determinant of a  $n$  order matrix defined as follows:

$$a_{i,k} = \begin{cases} 0 & \text{if } k > 1 + i \\ \binom{i+1}{k-1} & \text{otherwise} \end{cases}$$

This formula gives first Bernoulli numbers. If the factor  $(-1)^n$ , is omitted, since the only Bernoulli number from zero with odd index is  $B_1$ , we get the second Bernoulli numbers:

$$B_n = \frac{|A_n|}{(n+1)!}$$

Examples:

$$B_1 = \frac{|1|}{2!} = \frac{1}{2}$$

$$B_2 = \frac{\begin{vmatrix} 1 & 2 \\ 1 & 3 \end{vmatrix}}{3!} = \frac{1}{6}$$

$$B_3 = \frac{\begin{vmatrix} 1 & 2 & 0 \\ 1 & 3 & 3 \\ 1 & 4 & 6 \end{vmatrix}}{4!} = \frac{0}{24} = 0$$

$$B_4 = \frac{\begin{vmatrix} 1 & 2 & 0 & 0 \\ 1 & 3 & 3 & 0 \\ 1 & 4 & 6 & 4 \\ 1 & 5 & 10 & 10 \end{vmatrix}}{5!} = \frac{-4}{120} = -\frac{1}{30}$$

$$B_5 = \frac{\begin{vmatrix} 1 & 2 & 0 & 0 & 0 \\ 1 & 3 & 3 & 0 & 0 \\ 1 & 4 & 6 & 4 & 0 \\ 1 & 5 & 10 & 10 & 5 \\ 1 & 6 & 15 & 20 & 15 \end{vmatrix}}{6!} = \frac{0}{720} = 0$$



$$B_6 = \frac{\begin{vmatrix} 1 & 2 & 0 & 0 & 0 & 0 \\ 1 & 3 & 3 & 0 & 0 & 0 \\ 1 & 4 & 6 & 4 & 0 & 0 \\ 1 & 5 & 10 & 10 & 5 & 0 \\ 1 & 6 & 15 & 20 & 15 & 6 \\ 1 & 7 & 21 & 35 & 35 & 21 \end{vmatrix}}{7!} = \frac{120}{5040} = \frac{1}{42}$$

#### 4.1 First alternative: one matrix without the denominator (updating 15.9.2017)

For the properties of the determinants, the previous formula can be modified by deleting the denominator and by changing the elements of the matrix of  $n$  rows and  $n$  columns as follows:

$$a_{i,k} = \begin{cases} 0 & \text{if } k > 1 + i \\ \binom{i+1}{k-1} \frac{1}{i+1} & \text{otherwise} \end{cases}$$

Exemple:

$$B_6 = \begin{vmatrix} \frac{1}{2} & 1 & 0 & 0 & 0 & 0 \\ \frac{1}{3} & 1 & 1 & 0 & 0 & 0 \\ \frac{1}{4} & 1 & \frac{3}{2} & 1 & 0 & 0 \\ \frac{1}{5} & 1 & 2 & 2 & 1 & 0 \\ \frac{1}{6} & 1 & \frac{5}{2} & \frac{10}{3} & \frac{5}{2} & 1 \\ \frac{1}{7} & 1 & 3 & 5 & 5 & 3 \end{vmatrix} = \frac{1}{42}$$

#### 4.2 Second alternative with Von Staudt-Clausen theorem (updating 15.9.2017)

Similarly, taking into account the Von Staudt-Clausen theorem, we can compute separately numerator and denominator as integer numbers already reduced to the minimum terms. The numerator is:

$$num(B_n) = |A_n|$$

where  $|A_n|$  is the determinant of a  $n$  order matrix defined as follows:

$$a_{i,k} = \begin{cases} 0 & \text{if } k > 1 + i \\ \binom{i+1}{k-1} & \text{if } i+1 \text{ is prime and } i \text{ is a divisor of } n \\ \binom{i+1}{k-1} \frac{1}{i+1} & \text{otherwise} \end{cases}$$

Exemple:

$$num(B_6) = \begin{vmatrix} 1 & 2 & 0 & 0 & 0 & 0 \\ 1 & 3 & 3 & 0 & 0 & 0 \\ \frac{1}{4} & 1 & \frac{3}{2} & 1 & 0 & 0 \\ \frac{1}{5} & 1 & 2 & 2 & 1 & 0 \\ \frac{1}{6} & 1 & \frac{5}{2} & \frac{10}{3} & \frac{5}{2} & 1 \\ 1 & 7 & 21 & 35 & 35 & 21 \end{vmatrix} = 1$$

As is known, the denominator is easily calculated with the theorem of Von Staudt-Clausen. Therefore, it can also be obtained as a product of the elements of the second column of these matrices.

## 5.0 Theorem 1AU (updating 1.10.2017)

**It is proved that by dividing the polynomials to calculate the sum of  $n$  successive integer powers in the product between  $(n + 1)$  and another polynomial, the coefficients of the latter are obtained by reversing the sum between the unit matrix and the matrix alternate**

**signs extracted from the triangle of Tartaglia defined as in theorem 1A**

Applying the distributive property you have:

$$(U_m + \bar{A}_m) \begin{pmatrix} \sum_{i=1}^n i^1 \\ \sum_{i=1}^n i^2 \\ \sum_{i=1}^n i^3 \\ \dots \\ \sum_{i=1}^n i^m \end{pmatrix} = \begin{pmatrix} \sum_{i=1}^n i^1 \\ \sum_{i=1}^n i^2 \\ \sum_{i=1}^n i^3 \\ \dots \\ \sum_{i=1}^n i^m \end{pmatrix} + \bar{A}_m \begin{pmatrix} \sum_{i=1}^n i^1 \\ \sum_{i=1}^n i^2 \\ \sum_{i=1}^n i^3 \\ \dots \\ \sum_{i=1}^n i^m \end{pmatrix} =$$

and also:

$$= \begin{pmatrix} \sum_{i=1}^n i \\ \sum_{i=1}^n i^2 \\ \sum_{i=1}^n i^3 \\ \dots \\ \sum_{i=1}^n i^m \end{pmatrix} + \sum_{i=1}^n \bar{A}_m \begin{pmatrix} i \\ i^2 \\ i^3 \\ \dots \\ i^m \end{pmatrix} =$$

keeping in mind what is already shown in theorem 1A:

$$\sum_{k=1}^m \bar{a}_{j,k} i^k = i \sum_{k=1}^m \bar{a}_{j,k} i^{k-1} = i(i^j - (i-1)^j)$$

and then

$$= \begin{pmatrix} \sum_{i=1}^n i \\ \sum_{i=1}^n i^2 \\ \sum_{i=1}^n i^3 \\ \dots \\ \sum_{i=1}^n i^m \end{pmatrix} + \sum_{i=1}^n \begin{pmatrix} i(i - (i - 1)) \\ i(i^2 - (i - 1)^2) \\ i(i^3 - (i - 1)^3) \\ \dots \\ i(i^m - (i - 1)^m) \end{pmatrix} =$$

finally being

$$\begin{aligned} \sum_{i=1}^n i(i^m - (i - 1)^m) &= 1 \cdot 1^m + 2 \cdot 2^m - 2 \cdot 1^m + 3 \cdot 3^m - 3 \cdot 2^m + \dots \\ &\dots + (n - 1)(n - 1)^m - (n - 1)(n - 2)^m + nn^m - n(n - 1)^m = \\ &= -1^m - 2^m - 3^m - \dots - (n - 1)^m + nn^m = \\ &= -1^m - 2^m - 3^m - \dots - (n - 1)^m - n^m + n^m + nn^m = \\ &= (1 + n)n^m - \sum_{i=1}^n i^m \end{aligned}$$

For the transitive property of equality:

$$(U_m + \overline{A}_m) \begin{pmatrix} \sum_{i=1}^n i^1 \\ \sum_{i=1}^n i^2 \\ \sum_{i=1}^n i^3 \\ \dots \\ \sum_{i=1}^n i^m \end{pmatrix} = \begin{pmatrix} (n + 1)n \\ (n + 1)n^2 \\ (n + 1)n^3 \\ \dots \\ (n + 1)n^m \end{pmatrix}$$

Now, having the triangular matrix, the result of the sum, a determinant not null (m!), let's say

$$E_m = (\bar{A}_m + U_m)^{-1}$$

and multiply for this square matrix, on the left, the two members of the equation you get the theorem:

$$\begin{pmatrix} \sum_{i=1}^n i^1 \\ \sum_{i=1}^n i^2 \\ \sum_{i=1}^n i^3 \\ \dots \\ \sum_{i=1}^n i^m \end{pmatrix} = E_m \cdot \begin{pmatrix} (n+1)n \\ (n+1)n^2 \\ (n+1)n^3 \\ \dots \\ (n+1)n^m \end{pmatrix}$$

Example: case m=7

$$U_7 = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix} \quad \bar{A}_7 + U_7 = \begin{pmatrix} 2 & 0 & 0 & 0 & 0 & 0 & 0 \\ -1 & 3 & 0 & 0 & 0 & 0 & 0 \\ 1 & -3 & 4 & 0 & 0 & 0 & 0 \\ -1 & 4 & -6 & 5 & 0 & 0 & 0 \\ 1 & -5 & 10 & -10 & 6 & 0 & 0 \\ -1 & 6 & -15 & 20 & -5 & 7 & 0 \\ 1 & -7 & 21 & -35 & 35 & -21 & 8 \end{pmatrix}$$

from which

$$E_7 = (\bar{A}_7 + U_7)^{-1}$$

and for the demonstrated theorem

$$\begin{pmatrix} \sum_{i=1}^n i^1 \\ \sum_{i=1}^n i^2 \\ \sum_{i=1}^n i^3 \\ \sum_{i=1}^n i^4 \\ \sum_{i=1}^n i^5 \\ \sum_{i=1}^n i^6 \\ \sum_{i=1}^n i^7 \end{pmatrix} = \begin{pmatrix} \frac{1}{2} & 0 & 0 & 0 & 0 & 0 & 0 \\ \frac{1}{6} & \frac{1}{3} & 0 & 0 & 0 & 0 & 0 \\ 0 & \frac{1}{4} & \frac{1}{4} & 0 & 0 & 0 & 0 \\ -\frac{1}{30} & \frac{1}{30} & \frac{3}{10} & \frac{1}{5} & 0 & 0 & 0 \\ 0 & -\frac{1}{12} & \frac{1}{12} & \frac{1}{3} & \frac{1}{6} & 0 & 0 \\ \frac{1}{42} & -\frac{1}{42} & -\frac{1}{7} & \frac{1}{7} & \frac{5}{14} & \frac{1}{7} & 0 \\ 0 & \frac{1}{12} & -\frac{1}{12} & -\frac{5}{24} & \frac{5}{24} & \frac{3}{8} & \frac{1}{8} \end{pmatrix} \cdot \begin{pmatrix} (n+1)n \\ (n+1)n^2 \\ (n+1)n^3 \\ (n+1)n^4 \\ (n+1)n^5 \\ (n+1)n^6 \\ (n+1)n^7 \end{pmatrix}$$

which corresponds to

$$\begin{aligned} \sum_{k=1}^n k^1 &= (n+1) \frac{1}{2} n \\ \sum_{k=1}^n k^2 &= (n+1) \left( \frac{1}{6} n + \frac{1}{3} n^2 \right) \\ \sum_{k=1}^n k^3 &= (n+1) \left( \frac{1}{4} n^2 + \frac{1}{4} n^3 \right) \\ \sum_{k=1}^n k^4 &= (n+1) \left( -\frac{1}{30} n + \frac{1}{30} n^2 + \frac{3}{10} n^3 + \frac{1}{5} n^4 \right) \\ \sum_{k=1}^n k^5 &= (n+1) \left( -\frac{1}{12} n^2 + \frac{1}{12} n^3 + \frac{1}{3} n^4 + \frac{1}{6} n^5 \right) \\ \sum_{k=1}^n k^6 &= (n+1) \left( \frac{1}{42} n - \frac{1}{42} n^2 - \frac{1}{7} n^3 + \frac{1}{7} n^4 + \frac{5}{14} n^5 + \frac{1}{7} n^6 \right) \\ \sum_{k=1}^n k^7 &= (n+1) \left( \frac{1}{12} n^2 - \frac{1}{12} n^3 - \frac{5}{24} n^4 + \frac{5}{24} n^5 + \frac{3}{8} n^6 + \frac{1}{8} n^7 \right) \\ &\dots \dots \dots \end{aligned}$$

## 6.0 Theorem 1BU (updating 1.10.2017)

It is proved that by dividing the polynomials to calculate the sum of  $n$  successive integer powers in the product between  $(n - 1)$  and another polynomial, the coefficients of the latter are obtained by reversing the sum between the unit matrix and the matrix extracted from the triangle of Tartaglia defined as in theorem 1B

Applying the distributive property you have:

$$(U_m + A_m) \begin{pmatrix} \sum_{i=1}^{n-1} i^1 \\ \sum_{i=1}^{n-1} i^2 \\ \sum_{i=1}^{n-1} i^3 \\ \dots \\ \sum_{i=1}^{n-1} i^m \end{pmatrix} = \begin{pmatrix} \sum_{i=1}^{n-1} i^1 \\ \sum_{i=1}^{n-1} i^2 \\ \sum_{i=1}^{n-1} i^3 \\ \dots \\ \sum_{i=1}^{n-1} i^m \end{pmatrix} + A_m \begin{pmatrix} \sum_{i=1}^{n-1} i^1 \\ \sum_{i=1}^{n-1} i^2 \\ \sum_{i=1}^{n-1} i^3 \\ \dots \\ \sum_{i=1}^{n-1} i^m \end{pmatrix} =$$

and also:

$$= \begin{pmatrix} \sum_{i=1}^{n-1} i \\ \sum_{i=1}^{n-1} i^2 \\ \sum_{i=1}^{n-1} i^3 \\ \dots \\ \sum_{i=1}^{n-1} i^m \end{pmatrix} + \sum_{i=1}^{n-1} A_m \begin{pmatrix} i \\ i^2 \\ i^3 \\ \dots \\ i^m \end{pmatrix} =$$

keeping in mind what is already shown in theorem 1B:

$$\sum_{k=1}^m a_{j,k} i^k = i \sum_{i=1}^m a_{j,k} i^{k-1} = i((i+1)^j - i^j)$$

$$= \begin{pmatrix} \sum_{i=1}^{n-1} i \\ \sum_{i=1}^{n-1} i^2 \\ \sum_{i=1}^{n-1} i^3 \\ \dots \\ \sum_{i=1}^{n-1} i^m \end{pmatrix} + \sum_{i=1}^{n-1} \begin{pmatrix} i((i+1) - i) \\ i((i+1)^2 - i^2) \\ i((i+1)^3 - i^3) \\ \dots \\ i((i+1)^m - i^m) \end{pmatrix} =$$

and finally being

$$\begin{aligned} \sum_{i=1}^{n-1} i((i+1)^m - i^m) &= 1 \cdot 2^m - 1 \cdot 1^m + 2 \cdot 3^m - 2 \cdot 2^m + 3 \cdot 4^m - 3 \cdot 3^m + \dots \\ &\dots + (n-1)n^m - (n-1)(n-1)^m = \\ &= -1^m - 2^m - 3^m - \dots - (n-1)^m + (n-1)n^m = \\ &= (n-1)n^m - \sum_{i=1}^{n-1} i^m \end{aligned}$$

and then:

$$= \begin{pmatrix} \sum_{i=1}^{n-1} i \\ \sum_{i=1}^{n-1} i^2 \\ \sum_{i=1}^{n-1} i^3 \\ \dots \\ \sum_{i=1}^{n-1} i^m \end{pmatrix} + \begin{pmatrix} n(n-1) - \sum_{i=1}^{n-1} i \\ n^2(n-1) - \sum_{i=1}^{n-1} i^2 \\ n^3(n-1) - \sum_{i=1}^{n-1} i^3 \\ \dots \\ n^m(n-1) - \sum_{i=1}^{n-1} i^m \end{pmatrix} = \begin{pmatrix} (n-1)n \\ (n-1)n^2 \\ (n-1)n^3 \\ \dots \\ (n-1)n^m \end{pmatrix}$$



$$(U_m + A_m) \begin{pmatrix} \sum_{i=1}^{n-1} i^1 \\ \sum_{i=1}^{n-1} i^2 \\ \sum_{i=1}^{n-1} i^3 \\ \dots \\ \sum_{i=1}^{n-1} i^m \end{pmatrix} = \begin{pmatrix} (n-1)n \\ (n-1)n^2 \\ (n-1)n^3 \\ \dots \\ (n-1)n^m \end{pmatrix}$$

Now, having the triangular matrix, the result of the sum, a determinant not null (m!), let's say

$$\overline{E}_m = (A_m + U_m)^{-1}$$

and multiply for this square matrix, on the left, the two members of the equation you get the theorem:

$$\begin{pmatrix} \sum_{i=1}^{n-1} i^1 \\ \sum_{i=1}^{n-1} i^2 \\ \sum_{i=1}^{n-1} i^3 \\ \dots \\ \sum_{i=1}^{n-1} i^m \end{pmatrix} = \overline{E}_m \cdot \begin{pmatrix} (n-1)n \\ (n-1)n^2 \\ (n-1)n^3 \\ \dots \\ (n-1)n^m \end{pmatrix}$$

Example: case m=7

$$U_7 = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix} \quad A_7 + U_7 = \begin{pmatrix} 2 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 3 & 0 & 0 & 0 & 0 & 0 \\ 1 & 3 & 4 & 0 & 0 & 0 & 0 \\ 1 & 4 & 6 & 5 & 0 & 0 & 0 \\ 1 & 5 & 10 & 10 & 6 & 0 & 0 \\ 1 & 6 & 15 & 20 & 15 & 7 & 0 \\ 1 & 7 & 21 & 35 & 35 & 21 & 8 \end{pmatrix}$$

from which

$$\overline{E}_7 = (A_7 + U_7)^{-1}$$

and for the theorem demonstrated

$$\begin{pmatrix} \sum_{i=1}^{n-1} i^1 \\ \sum_{i=1}^{n-1} i^2 \\ \sum_{i=1}^{n-1} i^3 \\ \sum_{i=1}^{n-1} i^4 \\ \sum_{i=1}^{n-1} i^5 \\ \sum_{i=1}^{n-1} i^6 \\ \sum_{i=1}^{n-1} i^7 \end{pmatrix} = \begin{pmatrix} \frac{1}{2} & 0 & 0 & 0 & 0 & 0 & 0 \\ -\frac{1}{6} & \frac{1}{3} & 0 & 0 & 0 & 0 & 0 \\ 0 & -\frac{1}{4} & \frac{1}{4} & 0 & 0 & 0 & 0 \\ \frac{1}{30} & \frac{1}{30} & -\frac{3}{10} & \frac{1}{5} & 0 & 0 & 0 \\ 0 & +\frac{1}{12} & \frac{1}{12} & -\frac{1}{3} & \frac{1}{6} & 0 & 0 \\ -\frac{1}{42} & -\frac{1}{42} & +\frac{1}{7} & \frac{1}{7} & -\frac{5}{14} & \frac{1}{7} & 0 \\ 0 & -\frac{1}{12} & -\frac{1}{12} & +\frac{5}{24} & \frac{5}{24} & -\frac{3}{8} & \frac{1}{8} \end{pmatrix} \cdot \begin{pmatrix} (n-1)n \\ (n-1)n^2 \\ (n-1)n^3 \\ (n-1)n^4 \\ (n-1)n^5 \\ (n-1)n^6 \\ (n-1)n^7 \end{pmatrix}$$

which corresponds to

$$\sum_{k=1}^{n-1} k^1 = (n-1) \frac{1}{2} n$$

$$\sum_{k=1}^{n-1} k^2 = (n-1) \left( -\frac{1}{6} n + \frac{1}{3} n^2 \right)$$

$$\sum_{k=1}^{n-1} k^3 = (n-1) \left( -\frac{1}{4} n^2 + \frac{1}{4} n^3 \right)$$

$$\sum_{k=1}^{n-1} k^4 = (n-1) \left( \frac{1}{30} n + \frac{1}{30} n^2 - \frac{3}{10} n^3 + \frac{1}{5} n^4 \right)$$

$$\sum_{k=1}^{n-1} k^5 = (n-1) \left( \frac{1}{12} n^2 + \frac{1}{12} n^3 - \frac{1}{3} n^4 + \frac{1}{6} n^5 \right)$$

$$\sum_{k=1}^{n-1} k^6 = (n-1) \left( -\frac{1}{42} n - \frac{1}{42} n^2 + \frac{1}{7} n^3 + \frac{1}{7} n^4 - \frac{5}{14} n^5 + \frac{1}{7} n^6 \right)$$

$$\sum_{k=1}^{n-1} k^7 = (n-1) \left( -\frac{1}{12} n^2 - \frac{1}{12} n^3 + \frac{5}{24} n^4 + \frac{5}{24} n^5 - \frac{3}{8} n^6 + \frac{1}{8} n^7 \right)$$

## 7.0 References

(IT) Giorgio Pietrocola, [Esplorando un antico sentiero: teoremi sulla somma di potenze di interi successivi](#), Maecia, 2008.

(IT) Giorgio Pietrocola, [Internet e l'algoritmo di Ada Byron, contessa di Lovelace e incantatrice di numeri](#), Maecia, 2017

**(updating 23 giugno 2020)**

(IT) Giorgio Pietrocola [Sui polinomi per somme di potenze di interi successivi](#) 2018

In this other more recent work the theorems 1A and 1B are demonstrated simply by using vectors and Faulhaber's formula is proved and extended to any arithmetic progression.

(EN) Giorgio Pietrocola [From twin theorems on matrices obtainable from the Pascal triangle to the problem of sums of successive integers extended to any arithmetic progression](#) May-July 2019

(extracted and translated from Report presented at the 6th Mat ^ Nat symposium in Fontecchio)